

Charm Photoproduction in ep Collisions at HERA

Yehuda Eisenberg^{a*}

On behalf of the ZEUS Collaboration

^aWeizmann Institute of Science, Particle Physics Department, Rehovot, ISRAEL
email: yehuda@mail.desy.de

We report the latest results of the ZEUS collaboration on the photoproduction of D^* mesons in a low W range. Differential cross sections as function of $p_{\perp}^{D^*}$ and η^{D^*} are measured and compared with several recent NLO pQCD calculations. The differential cross-sections in a restricted kinematical region are higher than the NLO calculations, in particular in the forward (proton) direction. A recent pQCD model (BKL) describes the data reasonably well.

1. INTRODUCTION

Heavy quark photoproduction can be used to probe pQCD calculations with a hard scale given by the heavy quark mass and the high transverse momentum of the produced parton ($m_Q \gg \Lambda_{QCD}$). Two types of NLO calculations with different approaches are available for comparison with measurements of charm photoproduction at HERA. The massive charm approach [1] assumes light quarks to be the only active flavours within the structure functions of the proton and the photon, while the massless charm approach [2, 3] also treats charm as an active flavour and is thus only valid for $p_{\perp} \gg m_c$.

The data taken by the ZEUS collaboration during 1996/1997 corresponds to an integrated luminosity of about 37 pb^{-1} . In a subsample of about 17 pb^{-1} a small calorimeter positioned along the beam pipe was used to tag low W events, $80 < W_{\gamma p} < 120 \text{ GeV}$. The results of the high W region ($130 < W_{\gamma p} < 280 \text{ GeV}$) have been published before [4] and will not be shown here. This is the first presentation of our low W results. Charm was identified by the observation of $D^*(2010)$ mesons, which were reconstructed in the following decay modes: $D^{*+} \rightarrow D^0 \pi_s^+ \rightarrow (K^- \pi^+) \pi_s^+$ ($Br = 0.0262 \pm 0.0010$) and $D^{*+} \rightarrow D^0 \pi_s^+ \rightarrow (K^- \pi^+ \pi^+ \pi^-) \pi_s^+$ ($Br = 0.051 \pm 0.003$) and charge conjugates. The kinematic range studied was

$p_{\perp}^{D^*} > 2 \text{ GeV}$ and $-1.5 < \eta^{D^*} < 1.5$ for the high W region, and $2 < p_{\perp}^{D^*} < 8 \text{ GeV}$ and $-1.0 < \eta^{D^*} < 1.5$ for the low W region. The pseudorapidity is $\eta^{D^*} = -\ln(\tan \frac{\theta}{2})$, where θ is the polar angle with respect to the proton beam direction.

Charged tracks were measured in the central tracking detector. Cross sections were calculated in the photoproduction range of photon virtualities $Q^2 < 1 \text{ GeV}^2$ ($Q^2 < 0.015 \text{ GeV}^2$ for the tagged data).

2. D^* RECONSTRUCTION

D^* events have been selected by means of the mass difference (ΔM) method [5]. In the high W region we have observed [4] 3702 ± 136 D^* 's in the $D^0 \rightarrow (K\pi)$ decay mode with $p_{\perp}^{D^*} > 2 \text{ GeV}$, and 1397 ± 108 in the $(K\pi\pi\pi)$ decay mode with $p_{\perp}^{D^*} > 4 \text{ GeV}$ ($M(D^0) = 1.80\text{--}1.92 \text{ GeV}$). In the low W region we triggered only the $(K\pi)$ decay mode, and observed 550 ± 36 D^* events in the range $2 < p_{\perp}^{D^*} < 8 \text{ GeV}$ (Fig. 1). All tracks were assumed to be pions and kaons in turn; wrong charge D^* combinations [4] were used as a background distribution (dashed curve in Fig. 1), normalized outside the signal region.

3. D^* CROSS SECTIONS AND COMPARISON WITH CALCULATIONS

The D^* differential cross sections in the low W region are shown in Fig. 2 to 5. For comparing the

*Supported by the German Israeli Foundation and by the Minerva Foundation.




Figure 1. ΔM distribution for the events inside the D^0 mass region. The wrong charge distribution is shown as a dashed histogram.

experimental data to the NLO QCD calculations, we have used the D^* branching value measured by OPAL [6] $f(c \rightarrow D^{*+} + \dots) = 0.222 \pm 0.014 \pm 0.014$. For the charm fragmentation to D^* the Peterson [7] fragmentation function was used:

$$D_c(z) = N \frac{z(1-z)^2}{[(1-z)^2 + \epsilon z]^2}, \quad z = \frac{p_{D^*}}{p_c}.$$

In the massive calculation $\epsilon = 0.036$ was obtained from a recent fit of Nason and Oleari [8] to ARGUS data. Alternatively, the Peterson fragmentation was replaced by fragmentation effects estimated by a leading order Monte Carlo (Pythia). Initial and final state radiation were not included. The results of both calculations for the low W region are shown in Figs. 2 and 3. The cross sections are compared with NLO QCD massive calculations using MRSG for the proton structure function (SF) and GRV-G HO for the photon. The theoretical massive calculations are below the data, in particular in the forward (proton) direction, although the Pythia fragmentation slightly improves the agreement.

A comparison with massless calculations [2], which are expected to become valid mainly at higher $p_{\perp}^{D^*}$, is shown in Fig. 4 for several photon structure functions. Some sensitivity to the photon SF seems to be present, but the excess in




Figure 2. ZEUS tagged data compared to massive NLO calculations of $d\sigma/dp_{\perp}^{D^*}$.

the forward direction is evident. The structure function GS-G-96 HO describes the data best.

Recently Berezhnoy, Kiselev and Likhoded (BKL) have suggested a new model for describing D^* photoproduction [9]. In this tree level pQCD $O(\alpha_s^3)$ calculation, they hadronize the $(c, \bar{q})$ state produced in pQCD, taking into account higher twist terms at $p_{\perp} \simeq m_c$. Thus the model is supposed to be valid over the whole $p_{\perp}$ range studied. No explicit resolved component is used. Singlet and octet color states both contribute to D^* production. The color state ratio $O(8)/O(1)$ is a free parameter in this model and was tuned to the ZEUS untagged results [4], yielding a value of 1.3.

Comparison of these calculations, for the same Octet/Singlet mixture, with the ZEUS tagged low W data is shown in Fig. 5. A better agreement with the data is observed than that for the NLO calculations.




Figure 3. ZEUS differential cross sections $d\sigma/d\eta$ compared to the massive NLO predictions. The dashed/dashed-dotted lines correspond to the Pythia fragmentation. Parameter sets are as in Fig.2.




Figure 4. ZEUS data and massless NLO predictions [2] of differential cross sections for several photon structure functions. Peterson fragmentation was used with $\varepsilon = 0.116$.




Figure 5. Comparison of the ZEUS low W data with the BKL model. The Octet/Singlet ratio is 1.3, as tuned for the high W region.

REFERENCES

1. S. Frixione et al., Nucl. Phys. **B454** (1995) 3; Phys. Lett. **B348** (1995) 633.
2. B. Kniehl et al., Z. Phys. **C76** (1997) 689; J. Binnewies et al., Z. Phys. **C76** (1997) 677; Phys. Rev. **D58** (1998) 014014.
3. M. Cacciari et al., Phys. Rev. **D55** (1997) 2736, 7134.
4. J. Breitweg et al., ZEUS Collaboration, Eur. Phys. J. **C6** (1999) 67.
5. S. Nussinov, Phys. Rev. Lett. **35** (1975) 1672; G.J. Feldman et al., Phys. Rev. Lett. **38** (1977) 1313.
6. K.Ackerstaff et al., Eur. Phys. J. **C1** (1998) 439.
7. C. Peterson et al., Phys. Rev. **D27** (1983) 105.
8. P. Nason and C. Oleari, Phys. Lett. **B447** (1999) 327.
9. A.V. Berezhnnoy, V.V. Kiselev, and A.K. Likhoded, hep-ph/9901333.